

A Final Report for

HIGH ALTITUDE PLASMA INSTRUMENT (HAPI) DATA ANALYSIS

IN-46

17 CIT

14098

11P

Submitted to:

National Aeronautics and Space Administration
Attention: Ms. Cynthia L. Tart
Greenbelt, MD 20771

Submitted by:

J. L. Burch

(NASA-CR-189360) HIGH ALTITUDE
PLASMA INSTRUMENT (HAPI) DATA
ANALYSIS Final Report (Southwest
Research Inst.) 11 p

N94-36949


Unclass

SwRI Project 15-3320

Contract NAS5-33030

G3/46 0014098

APPROVED:



James L. Burch, Vice President
Instrumentation and Space Research Division



SOUTHWEST RESEARCH INSTITUTE
Instrumentation and Space Research Division
6220 Culebra Road, San Antonio, Texas 78238
(210) 684-5111 • FAX (210) 647-4325

TABLE OF CONTENTS

	PAGE
INTRODUCTION	1
I. SUMMARY OF ACCOMPLISHMENTS	2
A. Investigations of Source Mechanisms for Electron Conics	2
B. Investigations of Particle Acceleration Phenomena in Auroral Acceleration Regions	3
C. Observations of Birkeland Currents Throughout the Nightside Auroral Regions	4
D. Studies of the AKR Source Region	5
E. Plasma Injection Phenomena in the Polar Cusp	5
F. Support of Studies Led by Other DE Groups	6
II. PUBLICATIONS FROM CONTRACT NAS5-33030	7
III. OTHER REFERENCES	8

INTRODUCTION

The objectives of the Dynamics Explorer mission are to investigate the coupling of energy, mass and momentum among the Earth's magnetosphere, ionosphere and upper atmosphere. At launch, on August 3, 1981, DE-1 was placed into an elliptical polar orbit having an apogee of 23,130 km to allow global auroral imaging and crossings of auroral field lines at altitudes of several thousand kilometers. At the same time DE-2 was placed into a polar orbit, coplanar with that of DE-1 but with a perigee altitude low enough (309 km) for neutral measurements and an apogee altitude of 1012 km.

The DE-1 High Altitude Plasma Instrument (HAPI) provided data on low and medium energy electrons and ions from August 13, 1981 until December 1, 1981, when a high-voltage failure occurred. Analysis of the HAPI data for the time period of this contract has produced new results on: (1) the source mechanisms for electron conical distributions, (2) particle acceleration phenomena in auroral acceleration regions, (3) Birkeland currents throughout the nightside auroral regions, (4) the source region for auroral kilometric radiation (AKR), and (5) plasma injection phenomena in the polar cusp.

The DE Science Team, at its meeting on March 7 and 8, 1989, identified four Team Science Objectives:

1. Plasma physics of the auroral acceleration region,
2. Dynamical interactions between the thermosphere and the magnetosphere,
3. The ionosphere as a source of plasma for the magnetosphere,
4. Electrodynamic coupling as a function of substorm phase and IMF direction.

Most of the DE-1 HAPI research that has been done under this contract contributes to the first objective, plasma physics of the auroral acceleration region, although objectives (3) and (4) are also represented in the work.

I. SUMMARY OF ACCOMPLISHMENTS

Discussed below are some of the major accomplishments of the HAPI data analysis effort during the contract period.

A. *Investigations of Source Mechanisms for Electron Conics*

Electron conics were observed in the DE-1 data by Menietti and Burch [1985], who reported them to be associated with trapped particles and parallel electric fields. The authors suggested perpendicular heating via wave-particle interactions as a source

mechanism, in analogy with ion conic formation. Lundin et al. [1987] and Hultqvist et al. [1988] later reported observations of electron conics in the Viking data and suggested that a parallel potential difference that varies in magnitude over a fraction of an electron bounce period might explain electron conics that were observed to be associated with ion conics. For plasma parameters characteristic of those observed by DE-1 in the mid-altitude nightside auroral region, Wong et al. [1988] have shown that upper hybrid waves generated by a loss-cone distribution in a background cold plasma can preferentially heat the electron population in an magnetically oblique direction by cyclotron resonance. They pointed out that any particle distribution that produces $df/dv_{\perp} > 0$ might act as the free energy source.

Wong et al. [1988] also pointed out that because of the dependence of the diffusion coefficient on energy and pitch angle, upper hybrid waves will heat the electrons in both the parallel and perpendicular directions. By contrast, for the case of parallel acceleration only, the ratio of upgoing to downgoing electrons at any specific perpendicular energy should always be unity except in the case of lower energy backscattered electrons. For all of the observed electron conical distributions presented by Menietti and Burch [1985], however, more particles were detected moving up the field line than down, with the ratio as large as 1.45.

Recently Menietti et al. [1992] have reported observations by both particle and plasma wave instruments on board DE-1 and the Swedish Viking satellite that demonstrate the presence of intense upper hybrid waves at mid-altitudes of the polar magnetosphere in association with electron conical distributions. This work suggests the possibility that upper hybrid waves heat electrons perpendicular to the magnetic field and are responsible for at least some electron conical distributions. Of 12 DE-1 orbits on which electron conics were observed in the Menietti et al. [1992] study, waves near the upper hybrid frequency were seen (within about 30 seconds of the time of one frequency sweep of the plasma wave instrument, PWI) on 10 of them.

The above studies have thus far demonstrated by theory and simulation that electron conics can be generated by upper hybrid waves (perpendicular heating) although other sources such as, electron acoustic waves, Langmuir waves, and Alfvén-ion cyclotron waves have been suggested by others. More experimental and theoretical research remains to be done before the process or processes responsible for electron conical distributions can be identified with certainty.

B. *Investigations of Particle Acceleration Phenomena in Auroral Acceleration Regions*

Gurgiolo and Burch [1988] combined HAPI data and computer simulation techniques to determine the altitude profile of parallel potential distributions associated with auroral electron acceleration. An unexpected result was obtained. Agreement between the computer model and the HAPI data could only be found if the potential structure was split into a lower-altitude part and a higher-altitude part, with the spacecraft in a region containing no parallel electric fields. This was the first evidence of a split-type potential

distribution, and it can explain several other features of auroral ion and electron distributions.

The surprising result obtained by Gurgiolo and Burch [1988] raised the questions of what type of electron distribution would be expected to be observed within a region containing a parallel electric field and whether or not such distributions are ever seen in the HAPI data. The paper by Burch et al. [1990] reports the observation of a new type of auroral electron distribution, which has the characteristics of a bi-directional conic and which can be reproduced by test particle simulations. The conclusion of Burch et al. [1990] is that the typical midaltitude auroral electron distribution that has been observed by several different spacecraft is characteristic of a field-free region between parallel potential drops. The electron distribution that is expected within a parallel electric field is quite distinct and is only rarely observed by DE-1.

C. Observations of Birkeland Currents Throughout the Nightside Auroral Regions

By means of comparisons of the DE-1 HAPI data with the DE-1 MAG-A data, we have determined that at altitudes of several thousand to 20,000 kilometers upward-moving ionospheric electrons in regions of significant Birkeland currents are accelerated to energies above 5 eV, and hence all the major charge carriers can be observed by the HAPI instrument. This is not true at low altitudes, where upward-moving cold electrons, not observable with contemporary particle detectors, carry most, if not all, of the downward current. The major potential charge carriers are the accelerated cold electron beams (upward and downward), upward ion beams and conics, and precipitating electrons. While all of these populations, and sometimes others, can contribute significant current densities, we have found the cold electron beams to be the most common carriers of downward currents and, along with precipitating electrons, to carry the upward current as well.

In a recent published by Marshall et al. [1991] data from the DE-1 HAPI and MAG-A were used to examine particle acceleration and Birkeland current phenomena in the mid-altitude (10,000-20,000 km) nightside auroral region. Field-aligned current densities as derived from the two instruments were compared, generally resulting in good agreement. Comparisons were then made between the currents predicted by the model of the Knight [1973] for the parallel potential drops inferred from the HAPI data. It was found that large discrepancies resulted unless the full Knight formula (instead of the commonly used approximation) was used, including the temperatures of the observed electron distributions. The conclusion reached by Marshall et al. [1991] was that the Knight formula can predict fairly accurately the observed relationships between the parallel potential drop and the upward field-aligned current. Downward field-aligned currents, on the other hand, appear to be carried by suprathermal electron bursts, which may or may not be associated with parallel potential drops.

D. *Studies of the AKR Source Region*

According to Wu and Lee [1979] the most likely source of free energy for the generation of AKR is the loss-cone distribution within a region of depleted plasma density where the ratio of electron plasma frequency to gyrofrequency (ω_p/ω_g) is less than 0.3. This distribution provides the requirement of the cyclotron maser instability that $df/dv_{\perp} > 0$, and the theory has had success in explaining the observations to date.

Menietti et al. [1993] recently reported DE-1 particle and wave data for a pass of the nightside auroral region near an AKR source center. They compared their results to those reported by Viking and also to recent numerical simulation studies of Winglee and Pritchett [1986], which show a great similarity to the observations. As noted by Menietti et al. [1993], adjacent to and generally poleward of the AKR region was a region of electron distributions with $t_{\perp}/t_{\parallel} \gg 1$. Menietti et al. [1993] have presented DE-1 data that could indicate that a mechanism suggested by Winglee and Pritchett [1986] is responsible for producing a distribution with $df/dv_{\perp} > 0$ outside the forbidden region and that the cyclotron maser instability itself leads to population of the forbidden region as well as to the AKR wave growth.

The simulation studies of Winglee and Pritchett [1986] demonstrated how the competition between the bump-in-tail instability and the cyclotron maser instability can then produce diffusion along both the parallel and perpendicular directions, with the resulting distribution depending on the initial parameters such as the electron density and the wave propagation angle, but much more experimental evidence with data acquired at higher rates will be necessary to confirm this possibility.

E. *Plasma Injection Phenomena in the Polar Cusp*

The polar cusp is the only region of the magnetosphere through which solar-wind plasma has been observed to enter the magnetosphere directly. In addition to the direct penetration of solar-wind plasma, there is also propagation of waves that are excited at the magnetopause by processes such as the Kelvin-Helmholtz instability. Both phenomena carry energy down the polar cusp into the upper ionosphere where it can produce strong heating of ionospheric plasma. As a result of this heating, there is an enhanced outflow of ionospheric plasma known as the cleft ion fountain.

It has been suggested that downflowing ions in the cusp can generate waves around the lower hybrid frequency and multiples of the ion gyrofrequencies, and that such waves can cause ion conics. More recently Viking and DE-1 data have been used to study cyclotron resonance heating and double cyclotron absorption in the cusp. Along these same lines, Winglee et al. [1993a, b] have considered the injection of energetic magnetosheath plasma through the polar cusp and the ion heating and upward ion conics that may result. Their approach was to use Dynamics Explorer plasma measurements to characterize the two mesoscale phenomena that are easily observed in the cusp (the cusp ion flows and the large-scale ion conic population) and to explore the microscale processes responsible for the ion heating with particle simulations.

Observations from Dynamics Explorer 1 have shown that the injected magnetosheath ions have very distinct characteristics, forming a “V” shape on energy/pitch-angle spectrograms [Burch et al., 1982]. This unique spectral feature arises from the fact that the transit time from the injection region to the point of observation in the mid-altitude cusp is a function of ion pitch angle. Both impulsive injection and injection through a restricted region in a convection electric field can lead to the “V” signatures.

Winglee et al. [1993b] have recently shown from DE 1 data that cusp ion injection is subject to short period modulation (of about 10 - 18 seconds with an injection duration of only a few seconds). This modulation is shown to be associated with modulation of the characteristics of the outflowing ionospheric plasma owing to the interaction of magnetosheath and ionospheric plasma. There is clearly an association between these two mesoscale phenomena, which suggests that they are coupled by microscale processes occurring at temporal scales not observable with DE 1. Even with the available data, however, Winglee et al. [1993b] have noted a correlation between the cusp ion injections and the ion conics along with concurrent periodicities in upgoing and downgoing cusp electrons. The possibility that the magnetosheath ions can provide a net downward momentum transfer to the ambient ions as well as enhancing their heating rate was investigated by Winglee et al. [1993b] with electrostatic particle simulations.

The results of the simulations performed by Winglee et al. [1993b] are consistent with the DE-1 data in that the magnetosheath ion injections initially suppress the upward ion conics, as downward momentum is imparted to the ambient ions, but the conics strengthen as the ion injection diminishes as a result of a heating process, which is a slow speed ion-ion drift instability between the outflowing heavy ions and the H^+ ions.

F. Support of Studies Led by Other DE Groups

A number of other DE studies have been supported by our analysis of HAPI data for other groups. Several of these studies have been completed and are represented by publications listed in Section II.

II. PUBLICATIONS FROM CONTRACT NAS5-33030

- Burch, J. L., Diagnosis of Auroral Acceleration Mechanisms by Particle Measurements, in *Auroral Physics*, ed. by C.-I. Meng, M. Rycroft, and L. A. Frank, Cambridge University Press, 97-108, 1991.
- Burch, J. L., C. Gurgiolo, and J. D. Menietti, The Electron Signature of Parallel Electric Fields, *Geophys. Res. Lett.*, *17*, 2329-2332, 1990.
- Burch, J. L., J. D. Menietti, C. S. Lin, and R. L. Winglee, Mesoscale-Microscale Coupling Observed in Auroral and Cusp Region Plasmas, in press, Proceedings of AGU Chapman Conference on Micro and Meso Scale Phenomena in Space Plasmas, 1993.
- Burch, J. L., J. D. Menietti, and J. A. Slavin, Dayside Auroral Particle Acceleration Mechanisms Derived from Dynamics Explorer Data, *J. Geomag. and Geoelectr.*, *42*, 1365-1378, 1990.
- Burch, J. L., N. A. Saflekos, D. A. Gurnett, J. D. Craven, and L. A. Frank, The Quiet-Time Polar Cap: DE-1 Observations and Conceptual Model, *J. Geophys. Res.*, *97*, 19,403-19,412, 1992.
- Horwitz, J. L., C. J. Pollock, T. E. Moore, W. K. Peterson, J. L. Burch, J. K. Winningham, J. D. Craven, L. A. Frank, and A. Persoon, The Polar Cap Environment of Outflowing O⁺, *J. Geophys. Res.*, *97*, 8361-8379, 1992.
- Kainer, S., D. Krauss-Varban, J. D. Gaffey, Jr., C. S. Lin, and J. L. Burch, Stimulated Emission of AKR in Regions of a Weakly Unstable Electron Velocity Distribution, *J. Geophys. Res.*, *95*, 12,141-12,148, 1990.
- Lu, G., P. H. Reiff, and J. L. Burch, On the Auroral Current-Voltage Relationship, *J. Geophys. Res.*, *96*, 3523-3531, 1991.
- Marshall, J. A., J. L. Burch, J. R. Kan, P. H. Reiff, and J. A. Slavin, Sources of Field-Aligned Current in the Auroral Plasma, *Geophys. Res. Lett.*, *18*, 45-48, 1991.
- Menietti, J. D., and J. L. Burch, Particle Acceleration in the Auroral Region as Observed by DE-1, in press, Proceedings of AGU Chapman Conference on Auroral Plasma Dynamics, 1993.
- Menietti, J. D., and M. F. Smith, Inverted Vs Spanning the Cusp Boundary Layer, *J. Geophys. Res.*, *98*, 11,391-11,400, 1993.

- Menietti, J. D., J. L. Burch, R. M. Winglee, and D. A. Gurnett, DE-1 Particle and Wave Observations in Auroral Kilometric Radiation (AKR) Source Regions, *J. Geophys. Res.*, 98, 5865-5880, 1993.
- Menietti, J. K., C. S. Lin, H. K. Wong, A. Bahnsen, and D. A. Gurnett, Association of Electron Conical Distributions with Upper Hybrid Waves, *J. Geophys. Res.*, 97, 1353-1362, 1992.
- Reiff, P. H., G. Lu, J. L. Burch, J. D. Winningham, L. A. Frank, J. D. Craven, W. Peterson, and R. A. Heelis, On the High-and Low-Altitude Limits of the Auroral Electric Field Regions, in press, Proceedings of AGU Chapman Conference on Auroral Plasma Dynamics, 1993.
- Reiff, P. H., G. Lu, J. L. Burch, J. D. Winningham, L. A. Frank, J. D. Craven, W. K. Peterson, and R. A. Heelis, On the High-and Low-Altitude Limits of the Auroral Electric Field Region, in press, *J. Geophys. Res.*, 1993.
- Winglee, R. M., J. D. Menietti, W. K. Peterson, J. L. Burch, J. H. Waite, Jr., and B. Giles, Magnetosheath-Ionospheric Plasma Interactions in the Cusp/Cleft: 1. Observations of Modulated Injections and Upwelling Ion Fluxes, *J. Geophys. Res.*, 98, 19,315-19,330, 1993a.
- Winglee, R. M., J. D. Menietti, and C. S. Lin, Magnetosheath-Ionospheric Plasma Interactions in the Cusp/Cleft: 2. Mesoscale Particle Simulations, *J. Geophys. Res.*, 98, 19,331-19,348, 1993b.

III. OTHER REFERENCES

- Burch, J. L., P. H. Reiff, R. A. Heelis, J. D. Winningham, W. B. Hanson, C. Gurgiolo, J. D. Menietti, R. A. Hoffman, and J. N. Barfield, Plasma injection and transport in the mid-altitude polar cusp, *Geophys. Res. Lett.*, 9, 921-924, 1982.
- Gurgiolo, C., and J. L. Burch, Simulation of Electron Distributions within Auroral Acceleration Regions, *J. Geophys. Res.*, 93, 3989-4004, 1988.
- Hultqvist, B., R. Lundin, K. Stasiewicz, L. Block, P.-A. Lindqvist, G. Gustafsson, Koskinen, A. Bahnsen, T. A. Potemra, and L. J. Zanetti, Simultaneous observation of upward moving field aligned energetic electrons and ions on auroral zone field lines, *J. Geophys. Res.*, 93, 9765, 1988.
- Knight, S., Parallel electric fields, *Planet. Space Sci.*, 21, 741, 1973.
- Lundin, R., L. Eliasson, B. Hultqvist, and K. Stasiewicz, Plasma energization on auroral field lines as observed by the Viking satellite, *Geophys. Res. Lett.*, 14, 443, 1987.

- Menietti, J. D., and J. L. Burch, Electron conic signatures observed in the nightside auroral zone and over the polar cap, *J. Geophys. Res.*, *90*, 5345, 1985.
- Winglee, R. M., and P. L. Pritchett, The generation of low-frequency electrostatic waves in association with auroral kilometric radiation, *J. Geophys. Res.*, *91*, 13,531-13,541, 1986.
- Wong, H. K., J. D. Menietti, C. S. Lin, and J. L. Burch, Generation of electron conical distributions by upper hybrid waves in the Earth's polar region, *J. Geophys. Res.*, *93*, 10025, 1988.
- Wu, C. S., and L. C. Lee, A theory of the terrestrial kilometric radiation, *Astrophys. J.*, *230*, 621, 1979.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 1994	3. REPORT TYPE AND DATES COVERED Contractor Report	
4. TITLE AND SUBTITLE High Altitude Plasma Instrument (HAPI) Data Analysis			5. FUNDING NUMBERS 602	
6. AUTHOR(S) J. L. Burch				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Southwest Research Institute Instrumentation and Space Research Division 6220 Culebra Road San Antonio, TX 78238			8. PERFORMING ORGANIZATION REPORT NUMBER SwRI Project 15-3320	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, D.C. 20546-0001			10. SPONSORING/MONITORING AGENCY REPORT NUMBER CR-189360	
11. SUPPLEMENTARY NOTES Technical Monitor: G. Bullock/602				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 46 Report is available from the NASA Center for AeroSpace Information, 800 Elkridge Landing Road, Linthicum Heights, MD 21090; (301) 621-0390.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The objectives of the Dynamics Explorer mission are to investigate the coupling of energy, mass and momentum among the Earth's magnetosphere, ionosphere and upper atmosphere. At launch, on August 3, 1981, DE-1 was placed into an elliptical polar orbit having an apogee of 23,130 km to allow global auroral imaging and crossings of auroral field lines at altitudes of several thousand kilometers. At the same time DE-2 was placed into a polar orbit, coplanar with that of DE-1 but with a perigee altitude low enough (309 km) for neutral measurements and an apogee altitude of 1012 km.				
14. SUBJECT TERMS High Altitude Plasma Instrument, ionosphere, magnetosphere, thermosphere			15. NUMBER OF PAGES 9	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	